




ISSN: 1672 - 6553

JOURNAL OF DYNAMICS AND CONTROL

VOLUME 9 ISSUE 7: 183 - 204

SYNTHESIS, CHARACTERIZATION,
ANTICANCER AND BIOLOGICAL
ACTIVITIES OF TRANSITION METAL
COMPLEXES DERIVED OF
CHROMONE-3-CARBOXALDEHYDE
SCHIFF BASE LIGAND

Gulafsha D. Khan¹, Mahadeo A.
Sakhare¹, Sanjeevan A. Survase¹,
Yogesh N. Bharate¹, Shailendrasingh
V. Thakur^{2*}

¹Dept. of Chemistry, Balbhim Arts, Science &
Commerce College, Beed - 431122 (MS), India

²Dept. of Chemistry, Milliia Arts, Science &
Management Science College, Beed-431122
(MS), India

SYNTHESIS, CHARACTERIZATION, ANTICANCER AND BIOLOGICAL ACTIVITIES OF TRANSITION METAL COMPLEXES DERIVED OF CHROMONE-3-CARBOXALDEHYDE SCHIFF BASE LIGAND

Gulafsha D. Khan¹, Mahadeo A. Sakhare¹, Sanjeevan A. Survase¹, Yogesh N. Bharate¹, Shailendrasingh V. Thakur^{2*}

¹Dept. of Chemistry, Balbhim Arts, Science & Commerce College, Beed - 431122 (MS), India

²Dept. of Chemistry, Milliya Arts, Science & Management Science College, Beed-431122 (MS), India

*Corresponding Author: Shailendrasingh V. Thakur (khankg1994@gmail.com)

Abstract: A series of transition metal complexes of Co(II), Ni(II), Cu(II), Zn(II), Mn(II) and VO(II) have been prepared by tridentate Schiff base ligand containing N and O atoms. The Schiff base ligand(E)-1,5-dimethyl-4-(((4-oxo-4H-chromene-3-yl)methylene)amino)-2-phenyl-1H-pyrazol-3(2H)-one have been synthesized by reaction between chromone-3-carboxaldehyde and 4-aminoantipyrine and characterized by Mass, ¹H-NMR, FT-IR and UV-Visible spectroscopy and its metal complexes were characterised by FT-IR, UV-Visible, P-XRD, TGA and CHN analysis. All metal complexes exhibited 1:1 stoichiometry ratio of metal to ligand. The ligand and its metal complexes were screened for antibacterial, antifungal, anticancer and antioxidant activity. The biological activity of Schiff base ligand and its transition metal complexes has been studied by using four bacteria named as *Staphylococcus aureus*, *Bacillus subtilis*, *Salmonella typhi* and *Klebsiella pneumonia* and three fungi *Penicillium chrysogenum*, *Trichoderma viride*, *Aspergillus niger* by using disc diffusion method. The anticancer activity was screened by using MTT assay against human MCF-7 breast cancer cell line. The antioxidant activity was screened by using DPPH method. From all these observations it can be concluded that the Schiff base ligand and its transition metal complexes exhibited better activities.

Keywords: Schiff Base, chromone-3-carboxaldehyde, 4-aminoantipyrine, metal complexes, spectral and biological studies, etc.

1. Introduction:

During the last few decades special attentions given to the Schiff base transition metal complexes due to their structural diversity and as stereochemical models in coordination chemistry. Schiff base ligand and its transition metal complexes behaves as active anticancer drug.¹⁻² They are useful in some organic reaction and in magneto chemistry as a catalyst, it also useful in metalloproteins as models for metal containing sites in bioinorganic chemistry.³⁻⁴ The

Schiff base ligand synthesized from 4-aminoantipyrine and its transition metal complexes have various applications in clinical, analytical, pharmacological and biological fields.⁵⁻¹⁰ Schiff base prepared from 4-aminoantipyrine with aromatic aldehyde shows an important class of chelating ligand and metal complexes prepared are of great interest due to their great applications in industry, processes of transport, higher oxidation state stabilization, extraction and catalytic properties.¹¹⁻¹⁶ Such biological applications of these molecules motivate the researchers to prepare new compounds and to evaluate the biological activity by in vivo and in vitro studies. For spectrophotometric determination of metal ions antipyrine derivatives have been used. Most of these reagents gives several colours with transition metal ions which provides sensitive probes and some of the compound can also coordinate with rare earth metals to form interesting structures of metal complexes.¹⁷ Derivatives of pyrazolone are a group of antibiotics have been used in treatment of several bacterial diseases which have antibacterial activity against gram positive and gram negative strain.¹⁸ As per coordination point of view the ligands prepared from 4-aminoantipyrine has an advantage of two potential donor sites to prepare different types of compound with metal ions.¹⁹ In the present work Schiff base ligand and its transition metal complexes are prepared by condensation of 4-aminoantipyrine with chromone-3-carboxaldehyde and resulting transition metal complexes were investigated through different spectroscopic method and also these compounds are examined by antibacterial, antifungal, anticancer and antioxidant activity.

2. Experimental:

2.1 Materials methods:


In the present work all the chemicals used viz. chromone-3-carboxaldehyde, 4-aminoantipyrine and metal(II) acetate are of analytical grade and solvents like ethanol, ethyl acetate, petroleum ether, n-hexane which were used in the present work for synthesis are of spectroscopic grade. The microanalyses (C, H and N) were performed in 2400 Series II CHNS/O, Perkin Elmer analyser. The ¹H-NMR spectra of Schiff base ligand were recorded on BRUKER avance 400MHz ¹H-NMR instrument using DMSO as solvent. EI-MS spectra recorded on SYNAPT-XS#DBA06. The prepared Schiff base ligand and its metal complexes confirmed by FTIR spectra on JASCO FT/IR- 4700 spectrophotometer ranges from 4000 to 400 cm⁻¹. The electronic spectra recorded on UV-1800 series PERKIN ELMER UV-spectrophotometer by using DMSO as a solvent in between the range 200 to 1100 nm. Thermal analysis of the complexes performed under nitrogen atmosphere on TGA instruments

NETZSCH NJA-STA 2500 Regulus TGA thermal system. The Powder XRD was carried out on RIGAKU-JAPAN Miniflex 600 instrument.

2.2 Synthesis:

2.2.1 Synthesis of Schiff base ligand:


The Schiff base ligand (E)-1,5-dimethyl-4-(((4-oxo-4H-chromene-3-yl)methylene)amino)-2-phenyl-1H-pyrazol-3(2H)-one) was synthesized by condensation of ethanolic solution of chromone-3-carboxaldehyde (0.001 mol) and 4-aminoantipyrine (0.001 mol) by the literature method.²⁰ The reaction mixture was refluxed for 4-5 hours at 70-80°C and the reaction progress was monitored by thin-layer chromatography (TLC). The yellow solid product was formed, separated by filtration washed with cold ethanol and then dried under vacuum (Scheme1).




Scheme 1. Schematic route for the synthesis of Schiff base ligand

2.2.2 Synthesis of Schiff base transition metal(II) complexes (1-6):

A solution of metal acetate(II) in ethanol (0.001mol) was added to an ethanolic solution of Schiff base ligand (0.001mol) in 1:1 ratio and refluxed for 3-4 hours at 80-90°C. The pH of reaction mixture was maintained in between 7-8 by adding 10% alcoholic ammonia solution.²¹ The coloured solid products were obtained which were filtered, washed with cold ethanol and dried under vacuum.(Scheme 2)



M= Co(II), Ni(II), Cu(II), Zn(II) and Mn(II)



M = VO(II)

Scheme 2. Schematic route for the synthesis of Schiff base transition metal complexes (1-6).

2.3 Anticancer activity:

The Anticancer activity of Schiff base ligand and their metal complexes were carried out using MTT assay on MCF-7 human breast cancer cell line. MCF-7 human breast cancer cell line were used in DMEM medium with 10% fetal bovine serum and incubated in culture medium at a concentration of 1×10^4 cells/ml at 37°C and 5% CO_2 for 24h. Cells were put in (70 μl) 10^4 cells/well in 100 μl culture medium 100 μl sample (10, 40, 100 $\mu\text{g/ml}$) and control wells were incubated with DMSO and cell line. In triplicate all samples were incubated. Control were maintained to determine the survival of control cell and live cell percentage after culture and the cell cultures were incubated in 5% CO_2 incubator for 24 h at temperature of 37°C . After incubation, 20 μl of MTT reagent was added for complete removal of medium and then cells were incubated in CO_2 incubator for 4 hrs. Observed the wells through a microscope for the formation of formazan crystal. The yellow coloured MTT was reduced to dark coloured formazan only by viable cells. After the medium is completely removed then add 200 μl of DMSO (kept for 10 min) then incubated at 37°C . The absorbance of each sample was measured

against standard 5-fluorouracil using an Elisa micro plate reader (Benesphera E21) at a wavelength of 570 nm.^{22,23}

2.4 Antimicrobial activity:

The in vitro antimicrobial screening of Schiff base ligand and its metal complexes has been carried out against gram positive bacteria (*Staphylococcus aureus* and *Bacillus subtilis*) and gram negative bacteria (*Salmonella typhi* and *Klebsiella pneumonia*) and fungi (*Penicillium chrysogenum*, *Trichoderma viride* and *Aspergillus niger*) using disc diffusion method. The Muller-Hinton agar and Potato dextrose agar have been used as a medium for the growth of bacteria and fungi. All the compounds are dissolved in DMSO which is a negative control which shows no activity against bacteria and fungi. The sample plates were incubated for 24 h at 37°C. After 24 h the resulting inhibition zone were recorded in millimetre by using sliding calipers placed on the back side of the inverted petri plate. The minimum inhibitory concentration of Schiff base ligand and its transition metal complexes which showed antimicrobial activity was determined by using literature method.²⁴

2.5 Antioxidant assay:

The antioxidant activity assay of prepared Schiff base ligand and its transition metal complexes were tested by 1,1-diphenylpicrylhydrazyl (DPPH) method using Ascorbic acid as standard.²⁵ The standard solution was prepared by dissolving 1.95ml of 0.006% DPPH and 50µl of different concentrations (0-10 µg/50 µl) of standard ascorbic acid and taken the optical density of standard sample at 517nm. Then 10µl sample were dissolved in 40µl DMSO assed by 1.95ml 0.006% DPPH then % reduction in absorbance by ascorbic acid and also by sample is calculated. Then finally total antioxidant activity (TAA) in samples calculated by comparing it with standard ascorbic acid and expressed as mg.AAE/g sample (mg ascorbic acid equivalent per gram sample).

3. Result and discussion:

The Schiff base ligand and its Co(II), Ni(II), Cu(II), Zn(II), Mn(II) and VO(II) metal complexes are prepared and characterized by using different spectroscopic techniques like IR, ¹H NMR, Mass spectral analysis and also screened for biological activities. They are soluble in DMF and DMSO but insoluble in ethanol, methanol and water. The elemental analysis data of Schiff base and its metal complexes are good in accordance with the calculated values. Physical and analytical data of ligand and its metal complexes is shown in Table1.

Table 1. Physical and analytical data of Schiff base ligand and its transition metal complexes


Ligand/ Complexes	Colour	Yield (%)	Melting Point °C	Elemental analysis found (Calc) (%)		
				C	H	N
Ligand	Yellow	87	233	70.20(70.19)	4.40(4.73)	11.69(11.9)
[Co(II)L(H ₂ O)(oAC) ₂] 3H ₂ O	Brown	62	>300	48.18(49.30)	4.44(5.09)	7.55(6.90)
[Ni(II)L(H ₂ O)(oAC) ₂] 3H ₂ O	Dark Yellow	68	>300	48.07(49.37)	4.01(5.09)	7.01(6.90)
[Cu L(H ₂ O)(oAC) ₂] H ₂ O	Green	63	>300	52.04(51.98)	4.72(4.67)	7.28(7.27)
[Zn(II)L(H ₂ O)(oAC) ₂] 3H ₂ O	Orange	70	>300	48.83(48.78)	5.08 (5.04)	6.83 (6.84)
[Mn L(H ₂ O)(oAC) ₂] H ₂ O	Light Yellow	60	>300	52.82(52.77)	4.79 (4.74)	7.39 (7.38)
[VOL(H ₂ O)]SO ₄ H ₂ O	Black	59	>300	58.88(57.4)	4.47 (4.14)	9.81 (8.97)

3.1. IR spectra:

The IR spectral data of the synthesized Schiff base ligand and its metal complexes is compared to determine coordination sites involved in chelation. The IR spectra of Schiff base ligand and its transition metal complexes are shown in Figure.1. In the IR spectra of ligand, the band observed at 1651cm^{-1} is assigned to $\nu(\text{C}=\text{O})$ group of the chromone in free ligand on complexation it was shifted to lower frequency wave number indicates the coordination of carbonyl oxygen of the $\nu(\text{C}=\text{O})$ group to the metal ion.²⁶ Likewise C=O band at 1623cm^{-1} assigned to pyrazole carbonyl in free ligand which on complexation shifted to lower frequency wave number indicates that the participation of oxygen of pyrazole carbonyl is in chelation with the metal ion.²⁷ The C=N of (azomethine) shows band at 1567cm^{-1} in free ligand shifted to lower or higher frequency which indicates the participation of nitrogen of azomethine in coordination.²⁸ The IR spectral data of the metal complexes also shows new bands in the range of $590\text{-}535$ and $457\text{-}423\text{cm}^{-1}$ is due $\nu(\text{M-O})$ and $\nu(\text{M-N})$ respectively.²⁹ Therefore, from the IR spectral data it is concluded that the Schiff base ligand act as tridentate ligand with NOO donor sites which coordinate to the central metal ions via the C=O of the chromone, C=O of the pyrazole and nitrogen of azomethine. FT-IR data of Schiff base ligand and its metal complexes as shown in Table 2.

Table 2. FT-IR data of Schiff base ligand and its transition metal complexes

Sr. No.	Ligand/ Complexes	Chromone	Pyrazole	Azomethine	M-O	M-N
		C=O	C=O	C=N		
1.	L	1651	1623	1567	--	--
2.	Co(II)	1570	1570	1489	586	424
3.	Ni(II)	1610	1600	1555	584	440
4.	Cu(II)	1620	1598	1489	589	457
5.	Zn(II)	1605	1577	1489	582	454
6.	Mn(II)	1632	1590	1490	536	423
7.	VO(II)	1642	1610	1486	543	423

**Figure 1.** FT-IR Spectra of Schiff base ligand and transition metal complexes

3.2 ^1H -NMR spectra:

The ^1H -NMR spectral data of Schiff base ligand was taken at room temperature in DMSO solution. In the ^1H -NMR spectrum, singlet observed at 2.4δ and 3.2δ which attributes to the $-\text{CH}_3$, $-\text{NCH}_3$ group present in the pyrazole ring of Schiff base ligand. The ligand also shows the signal at 9.6δ attributed to singlet of azomethine proton ($-\text{CH}=\text{N}$).³⁰ and signal at 8.9δ gives singlet for the aromatic proton of chromone moiety.³¹ The bands at 7.4 - 8.1δ shows phenyl multiplet bands respectively. ^1H NMR (DMSO): δ = 2.4 (s, 3H, $-\text{CH}_3$), 3.2 (s, 3H, $-\text{NCH}_3$), 9.6 (s, 1H, $-\text{N}=\text{CH}$), 8.9 (s, $-\text{C}=\text{CH}$), 7.4 - 8.1 (m, 9H, Ar-H).




Figure 2. ^1H -NMR Spectra of Schiff base ligand

3.3 Mass spectra:

The molecular ion peaks verified the suggested formula for the synthesized compound. The mass spectra of Schiff base ligand exhibit the molecular ion peak at m/z 360 corresponds to $[\text{C}_{21}\text{H}_{17}\text{N}_3\text{O}_3]^+$ ion and 361 ($M+1$) which is in good agreement with the molecular weight of the proposed structure.




Figure 3. Mass Spectra of Schiff base ligand

3.4 TGA analysis:

The TGA analysis of transition metal complexes are taken which was recorded in the range of 25-800°C in nitrogen atmosphere with 30°C/10.0(K/min)/800°C heating rate and it can be observed that the transition metal complexes of Schiff base ligand decomposes at higher temperature range which concludes that metal complexes are formed. The Co(II), Ni(II), Cu(II), Zn(II), Mn(II) and VO(II) complexes of ligand were chosen for thermal study. Based on TGA data, the decomposition of all transition metal complexes occurs in two steps. In the first step, weight loss suggests the presence of lattice water and coordinated water molecules in Co(II), Ni(II), Cu(II), Zn(II) and Mn(II) complexes which are removed within the range of 50-150°C.³² It is observed from the TGA data that the Co(II), Ni(II) and Zn(II) complexes contain three lattice water molecules and one coordinated water molecule, while the Cu(II) and Mn(II) complexes contain one lattice water molecule and one coordinated water molecule. In the second step, the ligand is lost in the temperature range of 150– 750 °C along with the coordinated two acetate ions in Co(II), Ni(II), Cu(II), Zn(II) and Mn(II) complexes. In case of VO(II) complexes, In first step weight loss between temperature 50-250°C suggests the presence of one lattice water and one lattice sulphate ion and in second step weight loss between temperature 250-750°C corresponds to loss of ligand along with coordinate water molecule. The mass of final residue corresponds to stable metal oxide.³³ The TGA graph is shown in Figure 4.




Figure 4. TGA spectra of transition metal complexes

3.5 Electronic spectral analysis and molar conductivity:

The electronic spectra of Schiff base ligand and its metal complexes were recorded in DMSO in the range of 16666 to 50000 cm^{-1} and wavelength range of 200 to 1100 nm.³⁴⁻³⁸ The spectral data of Schiff base ligand and its metal complexes with molar conductance are demonstrate in Table 3. The molar conductivity were recorded in DMSO solvent and the values lies in between 2.1-9.8 $\text{cm}^2 \text{mol}^{-1}$ shows nonelectrolytic nature of Schiff base ligand and its transition metal complexes.^{39,40}

3.6 Powder x-ray diffraction:

The P-XRD of metal complexes is analysed in the range of $2\theta = 20-80^\circ$ at the wavelength of 1.540 Å which is useful for the knowledge of lattice parameters, cell parameters, crystal system. The pattern of diffraction shows the nature of transition metal complexes are crystalline.^{27,30} The P-XRD data of transition metal complexes is summarized in Table 4. Based on P-XRD data, the Ni(II) and Cu(II) complexes shows orthorhombic crystal system, Zn(II) and Mn(II) complexes shows monoclinic crystal system and Co(II) and VO(II) complexes shows triclinic crystal system.

Table 3. Electronic spectra of Schiff base ligand and its transition metal complexes

Ligand/ Complexes	Maximum absorption (nm)	Maximum absorption (cm ⁻¹)	Band assignment	Molar conductance S mol ⁻¹ cm ²
Ligand	268	37313	$n \rightarrow \pi^*$	2.1
	222	44943	$\pi \rightarrow \pi^*$	
Co(II)	402	24875	${}^4T_{1g}(F) \rightarrow {}^4T_{2g}(F)$	9.1
	328	30487	${}^4T_{1g}(F) \rightarrow {}^4A_{2g}(F)$	
	261	38240	${}^4T_{1g}(F) \rightarrow {}^4T_{1g}(P)$	
Ni(II)	371	26954	${}^3A_{2g}(F) \rightarrow {}^3T_{2g}(F)$	9.0
	279	35842	${}^3A_{2g}(F) \rightarrow {}^3T_{1g}(F)$	
	275	36363	${}^3A_{2g}(F) \rightarrow {}^3T_{1g}(P)$	
Cu(II)	338	29585	$2E_g \rightarrow {}^2T_{2g}$	7.2
	276	36231	Charge transfer	
Zn(II)	348	28694	Charge transfer	9.8
Mn(II)	385	25974	${}^3T_{1g}(F) \rightarrow {}^3T_{1g}(P)$	7.8
	390	25614	${}^3T_{1g}(F) \rightarrow {}^3A_{2g}(F)$	
	325	30721	Charge transfer	
	889	11248	${}^2B_2 \rightarrow {}^2E$	
	363	27548	Charge transfer	
VO(II)	276	36231	Charge transfer	8.1

Table 4. P- XRD data of transition metal complexes

Complexes	Co(II)	Ni(II)	Cu(II)	Zn(II)	Mn(II)	VO(II)
No. of reflection	17	32	17	33	30	35
Maxima (2 θ)	40.43	53.65	55.28	46.71	43.64	50.22
Intensity	100	100	100	100	100	100
d value	12.684	10.715	7.312	10.087	13.750	9.553
Lattice constant (Å)	a=7.765	a=7.150	a= 14.322	a= 8.393	a= 13.914	a =7.472
	b=10.59	b= 12.307	b= 13.320	b= 14.922	b= 7.500	b =8.235
	c=13.59	c= 21.431	c= 8.504	c= 10.093	c= 16.254	c =9.667
Unit cell volume	1014.274	1886.033	1622.300	1263.455	1676.434	570.875
Axis and axis angle	a $\neq$ b $\neq$ c	a $\neq$ b $\neq$ c	a $\neq$ b $\neq$ c	a $\neq$ b $\neq$ c	a $\neq$ b $\neq$ c	a $\neq$ b $\neq$ c
	and $\alpha \neq \beta \neq \gamma \neq 90^\circ$	and $\alpha = \beta = \gamma = 90^\circ$	and $\alpha = \beta = \gamma = 90^\circ$	and $\alpha = \gamma \neq \beta \neq 90^\circ$	and $\alpha = \gamma \neq \beta \neq 90^\circ$	and $\alpha \neq \beta \neq \gamma \neq 90^\circ$
Z Value	4	8	4	2	4	2
Crystal system	Triclinic	Orthorhombic	Orthorhombic	Monoclinic	Monoclinic	Triclinic



3.7 Anticancer activity:

The anticancer activity of synthesized Schiff base ligand and its metal complexes was measured in vitro at different concentration (10, 40, 100 µg/ml) using MTT assay method against MCF-7 human breast cancer cell line in comparison with standard 5-Fluorouracil shown in figure 5. It was observed that all the tested compound show good activity for different concentration some metal complexes shows lowest activity but are active in comparison with standard 5-Fluorouracil. Hence Schiff base ligand and its transition metal complexes are excellent anticancer agent against human breast cancer cell line MCF-7.^{26,41} Estimation based on IC₅₀ values of the transition metal complexes are higher than their parent Schiff base ligand which was compared with standard shows moderate values.⁴² Among these all the synthesized Schiff base ligand and its metal complexes show good activity as compared to standard compound. The % of inhibition for different concentration and IC₅₀ values of the prepared compound are presented in Table 5.

Table 5. Anticancer activity of Schiff base ligand and its transition metal complexes

Sr. No	Compounds	% of inhibition for Concentrations (µg/ml)			Ic ₅₀
		10	40	100	
1.	Ligand	72.77	74.3	74.19	24.56
2.	Co(II)	54.08	63.67	71.34	38.53
3.	Cu(II)	69.18	73.14	76.54	39.01
4.	Zn(II)	78.83	78.09	78.40	59.8
5.	5-Fluorouracil	79.14	83.29	88.79	42.08
	(Standard)				




Figure 5. Anticancer activity of Schiff base ligand and its transition metal complexes

3.8 Antimicrobial screening:

3.8.1 Antibacterial activity:

The antibacterial activity of prepared Schiff base ligand and its metal complexes were tested against *Staphylococcus aureus*, *Bacillus subtilis*, *Salmonella typhi* and *Klebsiella pneumoniae* by using disc diffusion method.⁴³ The results of antibacterial activity in graphical form are shown in figure 7. From findings it can be suggested that Schiff base ligand and its some metal complexes resistant up to 8 mm and sensitive in between 12-18 mm and highly sensitive ≥ 18 mm zone of inhibition against different bacteria. It was found that Schiff base ligand show higher activity than the corresponding transition metal complexes.⁴⁴ The zone of inhibition for prepared compounds are summarized in Table 6.




Figure 6. Antibacterial activity of Schiff base ligand and its transition metal complexes

Table 6. Antibacterial activity of Schiff base ligand and its transition metal complexes.

Compounds	Zone of inhibition(mm)			
	<i>S. aureus</i>	<i>B. subtilis</i>	<i>S. typhi</i>	<i>K. neumoniae</i>
Ligand	25.5	32	26.5	39.5
Co(II)	24.5	–	–	15
Ni(II)	16	–	–	18.5
Cu(II)	–	17	18.5	39
Zn(II)	–	23	17	38
Mn(II)	15	14.5	20	23.5
VO(II)	24	21.5	26.5	19
Control	-ve	-ve	-ve	-ve




Figure 7. Antibacterial activity of Schiff base ligand and its transition metal complexes

3.8.2 Antifungal activity:

The antifungal activity of prepared Schiff base ligand and its metal complexes were tested against *Penicillium chrysogenum*, *Trichoderma viride* and *Aspergillus niger* using disc diffusion method.⁴³ The findings in graphical form are shown in figure 9. From results it can be suggested that Schiff base ligand and its some metal complexes resistant to up to 8 mm and sensitive in between 12-18 mm and highly sensitive ≥ 18 mm zone of inhibition against

different fungi. Though the Schiff base ligand and its metal complexes possess good antifungal activity.⁴⁴ The zone of inhibition for of the prepared compounds are presented in Table 7.




Figure 8. Antibacterial activity of Schiff base ligand and its transition metal complexes

Table 7. Antifungal activity of Schiff base ligand and its transition metal complexes

Compounds	Zone of inhibition (mm)		
	<i>P. chrysogenum</i>	<i>T. viride</i>	<i>A. niger</i>
Ligand	17	20	14
Co(II)	15.5	20.5	24.5
Ni(II)	13.5	26.5	33
Cu(II)	25	24	–
Zn(II)	–	–	15.5
Mn(II)	12	12.5	19.5
VO(II)	25	28	31.5
Control	-ve	-ve	-ve




Figure 9. Antifungal activity of Schiff base ligand and its transition metal complexes

3.9 Antioxidant activity:

The antioxidants in the Schiff base ligand and their transition metal complexes were studied by DPPH method while Table.8 shows antioxidant values of prepared compounds at concentration of 10 μ l of sample compound and 40 μ l of DMSO are taken in which R denotes the reduction power. From finding it can be concluded that only zinc complex shows higher antioxidant activity in comparison with standard ascorbic acid.^{45,46}

Table 8. Antioxidant activity of Schiff base ligand and its metal complexes

Compounds	TAA on DPPH scale (mg.AAE/g)		
	R ₁	R ₂	R ₃
Ligand	-	-	-
Co(II)	-	-	-
Cu(II)	-	-	-
Zn(II)	23.38	23.23	22.69
Ascorbic acid (Standard)	19.79	19.70	19.99

4. Conclusion:

In this research paper we have synthesized the Schiff base ligand along with its transition metal complexes. Based on all spectroscopic techniques, ligand shows tridentate character. Based on P-XRD data, the Ni(II) and Cu(II) complexes shows orthorhombic crystal system, Zn(II) and Mn(II) complexes shows monoclinic and Co(II) and VO(II) complexes

show triclinic crystal system. From the biological studies of transition metal complexes, it is found that some transition metal complexes shows moderate antibacterial and antifungal activities than ligand and also some transition metal complexes shows good anticancer activities than ligand.

5. Acknowledgements:

The authors are gratefully acknowledging to the Principal Dr. Santosh S. Undare, Department of Chemistry, Balbhim Arts, Science and Commerce College, Beed (MS) for providing research facilities.

6. References:

1. Uddin N., Rashid F., Ali S., Tirmizi S A., Ahmad I., Zaib S., Zubair M., Diaconescu P. L., Tahir M. N., Iqbal J. and Haider A., "Synthesis, characterization, and anticancer activity of Schiff bases" *J. Biomol. Struct. Dyn.*, 38, 3246., (2020).
2. Mbugua S. N., Sibuyi N. R., Njenga L. W., Odhiambo R A., Wandiga S. O., Meyer M., Lalancette R. A. and Onani M. O., "New palladium(II) and platinum(II) complexes based on pyrrole Schiff bases: synthesis, characterization, x-ray structure and anticancer activity", *ACS omega* 5, 14942., (2020).
3. Kirkpatrick F. H., "Electrophoresis of large DNA molecules: theory and applications", *FAO. United Nations.*, (1990).
4. Raman N., Raja S. J., and Sakthivel A., "Transition metal complexes with Schiff base ligands: 4-aminoantipyrine based derivatives-a review", *J Coord Chem.*, 62, 691., (2009).
5. Donia A. M. and EI-Saled F. A., "Complexes of manganese(II), cobalt(II) and nickel(II) with the keto form of some antipyrine Schiff base derivatives", *Polyhedron* 7, 2149., (1998).
6. Joseph M., Nair K. M. and Radhakrishnan P. K., "Complexes of yttrium and lanthanide bromides with 4-N-(2'-hydroxybenzylidene) aminoantipyrine(1995)", *Synth React Inorg Met-Org Chem.*, 25, 1331., (1995).
7. Maurya R. C., Mishra D. D., Trivedi P. K., Mukherjee S. and Shrivastava P. K., "Synthesis and characterization of some novel penta coordinated mixed-ligand derivatives of zinc(II)-bis(acetylacetone) and bis(acetoacetanilide) chelates with 2/3-pyrazoline-5-one derivatives", *Synth React Inorg Met-Org Chem.*, 22, 403., (1992).

8. Agarwal R. K., and Singh G., "Preparation, spectral and thermal studies of some oxovanadium(IV) complexes of 4-aminoantipyrine", *Synth React Inorg Met-Org Chem.*, 16, 1183., (1986).
9. Chopra J. R., Uppal D., Arrora U. S, and Gupta S. K., "Synthesis and spectral studies of Cu(II) complexes of 4[N-(2-hydroxy-1-naphthalidene) amino] antipyrinethiosemicarbazones", *Asian J Chem.*, 12, 1277., (2000).
10. Radhakrishnan P. K., Indrasenan P, and Nair C. G. R., "Complexes of lanthanide nitrates with 4n-(2'-hydroxy benzylidene)-aminoantipyrine *Polyhedron* 3, 67., (1984).
11. Ismail K. Z., Dissouky A. E. and Shehada A. Z., "Spectroscopic and magnetic studies on some copper(II) complexes of antipyrine Schiff base derivatives", *Polyhedron* Vol. 16, 2909., (1984).
12. Singh L., Sharma A. and Sindhu S. K., "Structure and bonding of some newly synthesized complexes of lanthanide(III) complexes of 4[N-p-dimethylaminobenzalidene)amino]antipyrine", *Asian J. Chem.*, 11, 1445., (1999).
13. R. Bu. Xiu., F. L. Mintz., X. Z. You., Wang R. X., Yue Q., Meng Q. J., Lu Y. J. and Derveer D. V., "Synthesis and characteration of vanadyl complexes with unsymmetrical bis-schiff base ligands containing a cis-N₂O₂ coordinate chromophore", *Polyhedron*, 15 4885., (1996).
14. Chopra J. R., Uppal D., Arrora U. S. and Gupta S. K., "Synthesis and spectral studies of Cu(II) complexes of 4[N-2-hydroxy-1-naphthalidene)amino]antipyrine", *Asian J. Chem.*, 12, 1277., (2000).
15. Agarwal R. K., Garg P., Agarwal H. and Chandra S., "Synthesis, magneto-spectral and thermal studies of Co(II) and Ni(II) complexes of 4-[N-(N-(4-dimethylaminobenzylidene)amino)antipyrine", *Synth. React. Inorg. Met.-Org. Chem.*, 27, 251., (1997).
16. Singh L., Tyagi N., Dhaka N. and Sindhu S. K., "Synthesis, spectral and thermal studies of some lanthanide(III) complexes of 4-[N-(benzalidene)amino]antipyrinethiosemicarbazones", *Asian J. Chem.*, 11, 503., (1993).
17. Anupama B., Padmaja M. and Gyanakumari C., "Synthesis, characterization, biological activity and DNA binding studies of metal complexes with 4-aminoantipyrine Schiff base ligand", *E. J. Chem.*, 9, 389., (2012).
18. Raman N., Mitu L., Sakthivel A. and Pandi M. S. S., "Studies on DNA cleavage and antimicrobial screening of transition metal complexes of 4-aminoantipyrine derivatives of N₂O₂ type", *J. Iran. Chem. Soc.*, 6, 738., (2009).

19. Hossain M. E., Allam M. N., Begum J., Akbar A. M., Nazimuddin M., Smith F. E. and Hynes R. C., "The preparation characterization, crystal structure and biological activities of some copper(II) complexes of the 2-benzoylpyridine Schiff bases of S-methyl and S- benzyldithiocarbazate", *Inorg. Chim. Acta.*, 249, 207., **(1996)**.
20. Abdul khader Jailani N. M., Xavier A. and Ramu A., "Synthesis, spectroscopic characterization, DNA binding ability and biological activities of transition metal complexes containing tridentate Schiff base", *Materials Today: Proceedings.*, 5, 22200., **(2018)**.
21. Munde A. S., Jagdale A. N., Jadhav S. M. and Chondhekar T. K., "Synthesis, characterization and thermal study of some transition metal complexes of an asymmetrical tetradentate Schiff base ligand", *J. Serb. Chem. Soc.*, 75, 349., **(2010)**.
22. Sawant V. J., Bamane S. R., Shejwal R. V. and Patil S. B., "Comparision of drug delivery potentials of surface functionalized cobalt and zinc ferrite nanohybrids for curcumin in to MCF-7 breast cancer cells", *J. Magn. Magn. Mater.*, 417, 222., **(2016)**.
23. Manikyam H. K., Joshi S. K., Vakadi S. and Patil S. P., "Anticancer activity of novel terpenoidsaponin of psidiumguajava on MCF-7 cancer cell line using DAPI and MTT assays", *Afr. J. Pharm. Pharmacol.*, 15, 206., **(2021)**.
24. Shanker K., Rohini R., Ravinder V., Reddy P. M. and Ho. Y. P., "Ru(II) complexes of N₄ and N₂O₂ macrocyclic Schiff base ligands: their antibacterial and antifungal studies", *Spectrochimica Acta A: Molecular and Biomolecular Spectroscopy.*, 73, 205., **(2009)**.
25. Wong E., Chai T. and Hoo Y., "Antioxidation and cytotoxic activities of selected medicinal herbs used in Malaysia", *J. Med. Plantsres.*, 6, 3169., **(2012)**.
26. Kavitha P., Rama Chary M., Singavarapu B. V. V. A. and Reddy K. L., "Synthesis, characterization, biological activity and DNA cleavage studies of tridentate Schiff bases and their Co(II) complexes", *Journal of Saudi Chemical Society.*, **(2013)**.
27. Bharate Y. N., Sakhare M. A., Jadhav S. B.. and Naikwade S. D., "Synthesis, characterization and antibacterial screening of Mn(II), Co(II) acetate and VO(II) sulphate complexes of Schiff base ligand", *Rasayan J. Chem.*, 14, 479., **(2021)**.
28. Sece J. M., Quiros M. and Garmendina M. J. G., "Synthesis, X-ray crystal structure and spectroscopic, magnetic and EPR studies of copper(II) dimers with methoxy-di-(2-pyridyl)methoxide as bridging ligand", *Polyhedron* 19, 1005., **(2000)**.

29. Palakuri K. and Laxmareddy K., "Synthesis structural characterization, and biological activity studies of Ni(II) and Zn(II) Complexes", *Bioinorganic Chemistry and Applications.*, (2014).
30. Bharate Y. N., Sakhare M. A., Jadhav S. B. and Naikwade S. D., "Synthesis, characterization and antibacterial screening of Ni(II), Cu(II) and Zn(II) acetate complexes of Schiff base ligand", *Journal of Global Resource.*, 5, 345., (2019).
31. Mohamed G., El-Baradie K., El-Wakiel N. and Sara H., "Chromone Schiff base complexes: synthesis structural elucidation, molecular modeling, antitumor, antimicrobial, and DNA studies of Co(II), Ni(II), and Cu(II) complexes", *J. Iran. Chem. Soc.*, (2018).
32. Nikolaev A. V, Myachina L. I. and Logvinenko V.A., "Thermographic study of the structures of some simple, mixed and binuclear chelates of EDTA", *Therm AnalInorg Mater. and Phys. Chem.*, 2,779., (1969).
33. Aghao A. K., Janrao D. M., Janrao S. and Survase S. A., "Synthesis and characterization of Mn (II),Co(II), Ni(II), Cu(II) and Zn(II) complexes derived from 3-[(pyridine-2-ylimino) methyl] quinoxalin-2-ol", *Der Chemica Sinica*, 90, 5, (2015).
34. Al-Hamdani A. A. S., Balkhi A. M., Shaker S. A. and Falah A., "New azo-schiff base derived with Ni(II), Co(II), Cu(II), Pd(II) and Pt(II) complexes: Preparation, spectroscopic investigation, structural studies and biological activity" *J. Chil. Chem. Soc.*, 60 2774, (2015)
35. Ramaswamy P., Mandal S., Hegde N. N., Prabhu R., Banerjee D., Bhat S. V. and Natarajan S., "Synthesis, structure, and magnetic properties of amine-templated transition metal phosphate", *Eur. J. Inorg. Chem.*,1829, (2010).
36. Mishra A. P., Mishra R., Jain R. and Gupta S., "Synthesis of new VO(II), Co(II), Ni(II) and Cu(II) complexes with isatin-3-chloro-4-floroaniline and 2-pyridinecarboxylidene-4-aminoantipyrine and their antimicrobial studies", *The Korean Society of Mycology*, 40, 202012, (2015).
37. El-BoraeyHanaa A., Heba El-Ghnam, Ensaf M.. A. and Safaa S. H., "Designing of some VO(II), CO(II) and Hg(II) Schiff base complexes: Synthesis, structure elucidation, anticancer and biopesticidal activities, DFT and docking approaches", *Result in chem.*, 11, 101767., (2024).
38. Mendu P, GyanakumariC and Ragi R., "Synthesis, Characterization, DNA Binding, DNA Cleavage and Antimicrobial Studies of Schiff Base Ligand and its Metal Complexes" *J. Fluoresc.*, (2015).

39. Imran Ali, Wani W. A. and Kishwar S., “Empirical formulae to molecular structures of metal complexes by molar conductance”, *Syn Reac Inorg Met Org, Nano Metal Chem.* 43, 1162., **(2013)**.
40. Kurdekar G S, Sathisha M. P, Budagumpi S, Kulkarni N. V, Revankar V. and Suresh D. K., “4-Aminoantipyrine-based Schiff-base transition metal complexes as potent anticonvulsant agents”, *Med. Chem. Res.*, 21, 227., **(2012)**.
41. Zhong-Ying Ma, XinQiao, Cheng-ZhiXie, Jia Shao, Jing-Yuan Xu, Zhao-Yan Qiang and Jian-Shi Lou., “Activities of a novel Schiff Base copper(II) complex on growth inhibition and apoptosis induction toward MCF-7 human breast cancer cells via mitochondrial pathway”, *Journal of Inorganic Biochemistry*, 1171-19., **(2012)**.
42. Maria Sadia, Jehangir Khan, RobinaNaz, Muhammad Zahoor , Syed W. A. S , Riaz Ullah, Sumaira Naz, Ahmed Bari, Hafiz Majid Mahmood, Syed S. A., Ansari S. A. and Muhammad Sohaib., “Schiff base ligand L synthesis and its evaluation as anticancer and antidepressant agent”, *Journal of King Saud University-Science.*, 33, 101331, **(2021)**.
43. Survase S. A., Sakhare M. A., Bharate Y. N. and Patil C. S., “Synthesis, characterization and Biological investigation of Ni(II), Cu(II), Co(II), Zn(II), Mn(II) metal complexes of tetradentate Schiff base ligand”, *Journal of dynamics and control*, vol. 8, 9., **(2024)**.
44. Sarwade K. N., Sakhare M. A, Sakhare K. B. and Thakur S. V., Synthesis of metal-based biologically active agents from ONO-donor Schiff base ligand *Mongolian Journal of Chemistry* 25(52), **(2024)**, xx-xx
45. Shakir M, Summaiya Hanif, Owais Mohammad, Mohd. Asif Sherwani and Saud I. Al-Resayes., “Pharmacologically significant complexes of Mn(II), Co(II), Ni(II), Cu(II) and Zn(II) of novel Schiff base ligand, (E)-N-(furan-2-yl methylene) quinolin-8-amine: Synthesis, spectral, XRD, SEM, antimicrobial, antioxidant and in vitro cytotoxic studies”., *Journal of Molecular Structure.*, 1092, 143, **(2015)**.
46. Abdel Saeed A. M, Eishah M. A. M. and Abdul-Rahman Alawi Bin Yahia., “Synthesis, characterization and bioactivity evaluation of several divalent transition metal ion-bioflavonoid complexes”, *Results in chemistry.*, 7, 101535, **(2024)**.